arXiv:cond-mat/0608053v1 [cond-mat.other] 2 Aug 2006

Global singularity-free solution of the Iordanskii force problem

Markus Flaig and Uwe R. Fischer
Eberhard-Karls-Universität Tübingen, Institut für Theoretische Physik
Auf der Morgenstelle 14, D-72076 Tübingen, Germany
(Dated: February 6, 2008)

We present a derivation of the transverse force acting on a hydrodynamic vortex in the presence of an incoming sound wave from a global solution of the scattering problem, using the method of matched asymptotic expansions. The solution presented includes a detailed treatment of the interaction of the incident wave with the vortex core, and is free from the singularities in the momentum exchange between vortex and sound wave which have led to contradictory results for the value of the transverse force in the literature.

I. INTRODUCTION

The transverse force on a quantized vortex due to phonon scattering, first found by Iordanskii [1], has been controversially discussed in particular since the work of Thouless, Ao and Niu, which predicted that it in fact would vanish [2, 3, 4, 5, 6, 7, 8, 9, 10, 11]. Based on general properties of *superfluid* order and the Berry phase associated with the vortex motion, they came to the conclusion that the total transverse force is proportional to the superfluid density alone and given solely by the superfluid Magnus force, and not, as the existence of the Iordanskii force would imply, proportional to the *total density*, normal plus superfluid. Recent calculations based on the vector potential due to the Berry phase of quasiparticles lead to the same conclusion of a vanishing Iordanskii force [11].

It therefore seems in order to reinvestigate the assumptions underlying calculations of the Iordanskii force based on classical (but two-fluid) hydrodynamics (see, e.g., [4, 8]), where the transverse force stems from the momentum transfer between asymmetrically scattered sound waves and the vortex [12, 13]. The existing treatments use various simplifications for a treatment of this problem, which constitute sources for possible ambiguities and problems in the final result obtained for the value of the transverse force. First of all, the authors usually consider point vortices. However, for small distances to the vortex axis, the corresponding velocity field cannot be taken literally since then the fluid velocity would diverge and the Mach number of the problem would become infinite. It is not *a priori* clear that replacing the point vortex by a realistic vortex core would not affect the oscillatory motion of the vortex core due to the incident wave, and thus the momentum transfer between vortex and sound wave. Second, using simplified scalar wave equations instead of the full linearized hydrodynamic equations, one does not take into account the effects due the interaction of the incident wave with the vortex core leading to its oscillatory motion accurately [4]. Finally, working, as is conventional, with an (unphysical) infinitely extended plane wave results in mathematical difficulties (due to the non-convergence of sums over partial wave shifts in the scattering cross section [5]). We will show that no such singularities appear in the momentum transfer between sound wave and vortex, provided one is using a realistically shaped incident wave of a general form.

In the following, we present a study of the scattering problem which avoids using the simplifications discussed in the above. In particular, we derive a global solution for the coupled hydrodynamic equations using the method of matched asymptotic expansions, matching the solution for the vortical region, with a realistic core of finite extension, smoothly to the far-field wave region solution, assuming a general incoming wave instead of a plane wave. As a result, we obtain a regular expression for the transverse momentum exchange between sound field and vortex. When interpreted in terms of two-fluid hydrodynamics, this momentum exchange exactly corresponds to the Iordanskii force.

II. BASIC HYDRODYNAMIC EQUATIONS

Our starting point are the Euler and continuity equations for an ideal fluid described in terms of density ρ, velocity $\boldsymbol{v}$ and barotropic pressure functional $P = P(\rho)$:

$$\frac{\partial \rho}{\partial t} + \boldsymbol{\nabla} \cdot (\rho \boldsymbol{v}) = 0, \tag{1a}$$

$$\frac{\partial \boldsymbol{v}}{\partial t} + (\boldsymbol{v} \cdot \boldsymbol{\nabla}) \boldsymbol{v} = -\frac{1}{\rho} \boldsymbol{\nabla} P. \tag{1b}$$

In order to consider small perturbations $(\rho_1, \boldsymbol{v}_1)$ on a given background solution $(\rho_0, \boldsymbol{v}_0)$, which we require to satisfy the hydrodynamic equations, we introduce the following expansions of velocity and density:

$$\boldsymbol{v} = \boldsymbol{v}_0 + \boldsymbol{v}_1, \qquad \rho = \rho_0 + \rho_1. \tag{2}$$

Inserting these ansätze into (1), and equating terms of first order on both sides results in the following equations:

$$\frac{\mathrm{d}\rho_1}{\mathrm{d}t} + \boldsymbol{\nabla} \cdot (\rho_0 \boldsymbol{v}_1) + \rho_1 \boldsymbol{\nabla} \cdot \boldsymbol{v}_0 = 0, \tag{3a}$$

$$\frac{\mathrm{d}\boldsymbol{v}_1}{\mathrm{d}t} + (\boldsymbol{v}_1 \cdot \boldsymbol{\nabla}) \boldsymbol{v}_0 = -\boldsymbol{\nabla} \left(\frac{c^2}{\rho_0} \rho_1 \right); \tag{3b}$$

where the sound speed c is as usual defined by $c^2 = \partial P/\partial\rho$ and $\mathrm{d}/\mathrm{d}t = (\partial/\partial t + \boldsymbol{v}_0\cdot\boldsymbol{\nabla})$ denotes the convective derivative. We now switch from $(\rho_1, \boldsymbol{v}_1)$ to another set of variables $(\psi, \boldsymbol{\xi})$ via the following decomposition of the perturbations [15]

$$\begin{aligned}\rho_1 &= -\frac{\rho_0}{c^2}\frac{\mathrm{d}\psi}{\mathrm{d}t},\\ \boldsymbol{v}_1 &= \boldsymbol{\nabla}\psi + \boldsymbol{\xi}.\end{aligned} \tag{4}$$

The velocity perturbation is decomposed into a part with zero vorticity, $\boldsymbol{\nabla}\psi$, where the *pressure potential* ψ is regular, and a part comprising the vorticity in the sound wave, $\boldsymbol{\xi}$.

Our linearized hydrodynamic equations (3) may then be rewritten as

$$\frac{\mathrm{d}}{\mathrm{d}t}\left(\frac{1}{c^2}\frac{\mathrm{d}\psi}{\mathrm{d}t}\right) = \frac{1}{\rho_0}\boldsymbol{\nabla}\cdot[\rho_0(\boldsymbol{\nabla}\psi + \boldsymbol{\xi})], \tag{5a}$$

$$\frac{\mathrm{d}\boldsymbol{\xi}}{\mathrm{d}t} + (\boldsymbol{\xi}\cdot\boldsymbol{\nabla})\boldsymbol{v}_0 - \boldsymbol{\nabla}\psi\times\boldsymbol{\omega}_0 = 0. \tag{5b}$$

The above coupled pair of equations has first been derived by Bergliaffa et. al. using Clebsch potentials [15]. Eq. (5a) with the vortical part of the fluid velocity perturbations, $\boldsymbol{\xi}$, set to zero, was derived in [16, 17]. It can be shown [15], that in the case where the frequency Ω of the sound wave is much greater than the background vorticity $\boldsymbol{\omega}_0 = \boldsymbol{\nabla}\times\boldsymbol{v}_0$, there exist solutions where $\boldsymbol{\xi}$ is smaller by a factor $\boldsymbol{\omega}_0/\Omega$ than ψ. This leaves us with only one equation for the determination of the scalar quantity ψ when the *quasiclassical* condition $\omega_0/\Omega \ll 1$ is fulfilled; see the analysis of the *wave region* in the section to follow.

We study an axisymmetric vortex, which may be described by the following ansatz for the background solution, employing a cylindrical system of coordinates (r, ϕ, z) in what follows:

$$\rho_0 = \rho_0(r), \qquad \boldsymbol{v}_0 = v_0(r)\,\boldsymbol{e}_\phi. \tag{6}$$

The radial velocity v_0 is linked to the vorticity $\boldsymbol{\omega}_0 = \omega_0\,\boldsymbol{e}_z$ (which we consider to be a fixed quantity) by the relation

$$v_0(r) = \frac{1}{r}\int_0^r r'\omega_0(r')\,\mathrm{d}r'. \tag{7}$$

We require the vorticity for large r to decay faster than any power of r. This will later on allow us to use an approach based on the method of matched asymptotic expansions [18, 19]. For the velocity this assumption implies that, for $r\to\infty$, it is given by $v_0 = \gamma/r$ (here and below we ignore exponentially small terms), with $\gamma = \int_0^\infty r'\omega_0(r')\,\mathrm{d}r'$, which is the circulation divided by 2π. While the continuity equation (1a) is automatically satisfied by the ansatz (6), the density ρ_0 must be determined such that the Euler equation (1b) be fullfilled, which then leads to the following equation:

$$\rho_0' = \frac{\rho_0}{c^2}\frac{v_0^2}{r}, \tag{8}$$

where the prime denotes derivation with respect to r. We immediately see that density variations are of second order with respect to the Mach number $\mathscr{M}$ which is defined as $\mathscr{M} = \mathscr{U}/c_\infty$, with $\mathscr{U}$ being some typical velocity in the vortex and $c_\infty = c(r\to\infty)$ the sound speed for the fluid at rest at infinity. We require the Mach number to be a small quantity ($\mathscr{M} \ll 1$), which allows for an expansion of the solution of the hydrodynamic equations in $\mathscr{M}$ in the section to follow.

III. MATCHED ASYMPTOTIC EXPANSIONS

To carry out the matching procedure, we divide the plane into two regions: An inner, *vortical* region characterized by $r \simeq$, in which the vorticity is concentrated, and an outer, *wave* region, where r is of the order of the typical wavelength $\lambda = 2\pi\Omega/c_\infty$ of the incoming wave, which we take within the matched asymptotic expansion procedure to be $\lambda \simeq \mathscr{M}^{-1}$. After having developed suitably scaled non-dimensional equations for each region, the solutions in both region are expanded in the Mach number and matched where the regions overlap.

In the vortical region we rescale the spatial variable $\boldsymbol{r}$ according to $\boldsymbol{r}\to\boldsymbol{r}$ and the velocity $\boldsymbol{v}_0$ according to $\boldsymbol{v}_0 \to \mathscr{U}\boldsymbol{v}_0$. The typical time scale of the problem is given by the inverse of the frequency $\Omega^{-1} \simeq /\mathscr{U}$. Consequently, the appropriate scaling for the time variable t is $t \to (/\mathscr{U})t$, and the frequency Ω takes the scaled form $(\mathscr{U}/)\Omega$. Thus we have $|\boldsymbol{\omega}_0|/\Omega = \mathcal{O}(1)$, which means, in the light of the remarks in the previous section, that we should replace $\boldsymbol{\xi}$ by $\boldsymbol{\xi}/$. With these scalings, and using Eq. (8), our Eq. (5a) becomes:

$$\boldsymbol{\nabla}\cdot(\boldsymbol{\nabla}\psi + \boldsymbol{\xi}) = \mathscr{M}^2\left[\frac{\mathrm{d}^2\psi}{\mathrm{d}t^2} - \frac{v_0^2}{r}(\partial_r\psi + \xi_r)\right] + \mathcal{O}(\mathscr{M}^4), \tag{9}$$

while Eq. (5b) stays formally the same.

In the wave region, the appropriately scaled spatial variable is given by $\boldsymbol{R} = \mathscr{M}\boldsymbol{r}$. Furthermore, we define the velocity potential Ψ in the wave region as $\Psi(\boldsymbol{R}) = \psi(\boldsymbol{r})$. Since in the wave region the velocity is one order in the Mach number smaller than in the vortical region, we introduce the wave-region velocity $\boldsymbol{V}_0 = \mathscr{M}^{-1}\boldsymbol{v}_0 = \frac{\gamma}{R}\,\boldsymbol{e}_\phi$. Now, in the wave region, Eq. (9) takes on the following form:

$$(\boldsymbol{\nabla}^2 - \partial_t^2)\Psi = 2\mathscr{M}^2(\boldsymbol{V}_0\cdot\boldsymbol{\nabla})\partial_t\Psi + \mathcal{O}(\mathscr{M}^4). \tag{10}$$

Here and in what follows, we use the convention that the nabla operator acting on a wave-region quantity corresponds to a derivative with respect to $\boldsymbol{R}$. The equation for $\boldsymbol{\xi}$ is not needed because we restrict ourselves to solutions where in the wave region the vortical part of the velocity perturbation, $\boldsymbol{\xi}$, is exponentially small.

In order to decouple the equations in the vortical region, we split $\boldsymbol{\xi}$ into an irrotational and a divergence free part according to

$$\boldsymbol{\xi} = \boldsymbol{\nabla}\eta + \boldsymbol{e}_z\times\boldsymbol{\nabla}\zeta, \tag{11}$$

where η and ζ are uniquely specified by demanding that $\eta, \zeta \to 0$ as $r \to \infty$. Our Eq. (9) then becomes

$$\begin{aligned}\boldsymbol{\nabla}^2(\psi+\eta) &= \mathscr{M}^2\left(-i\varOmega+\boldsymbol{v}_0\cdot\boldsymbol{\nabla}\right)^2\psi \\ &\quad -\mathscr{M}^2\,\frac{v_0^2}{r}\left[\partial_r(\psi+\eta)-r^{-1}\partial_\phi\zeta\right]+\mathcal{O}(\mathscr{M}^4). \qquad (12)\end{aligned}$$

This Poisson equation may be solved with standard methods. Using a relation following from Eq. (5b)

$$\frac{\partial\boldsymbol{\xi}}{\partial t}-\boldsymbol{v}_0\times(\boldsymbol{\nabla}\times\boldsymbol{\xi})+\boldsymbol{\nabla}(\boldsymbol{v}_0\cdot\boldsymbol{\xi})-(\boldsymbol{\nabla}\psi+\boldsymbol{\xi})\times\boldsymbol{\omega}_0=0, \qquad (13)$$

and operating with the curl, we arrive at the following equation:

$$\begin{aligned}&\left(-i\varOmega+\frac{v_0}{r}\frac{\partial}{\partial\phi}\right)\boldsymbol{\nabla}^2\zeta-\frac{\omega_0'}{r}\frac{\partial\zeta}{\partial\phi} \\ &\qquad = -\omega_0'\frac{\partial}{\partial r}(\psi+\eta)-\omega_0\boldsymbol{\nabla}^2(\psi+\eta). \qquad (14)\end{aligned}$$

Note that here the fact that $\boldsymbol{\nabla}\cdot\boldsymbol{v}_0=0$ has been used. The above equation allows us to calculate ζ after $\psi+\eta$ has been calculated using (12).

We now formally expand the general solution in the Mach number $\mathscr{M}$ and then solve the resulting equations to the relevant order in the Mach number. In the vortical region the expansion takes on the following form

$$\begin{aligned}\psi &= \psi^{(0)}+\mathscr{M}\psi^{(1)}+\mathscr{M}^2\psi^{(2)}+\dots, \\ \eta &= \eta^{(0)}+\mathscr{M}\eta^{(1)}+\mathscr{M}^2\eta^{(2)}+\dots, \qquad (15) \\ \zeta &= \zeta^{(0)}+\mathscr{M}\zeta^{(1)}+\mathscr{M}^2\zeta^{(2)}+\dots;\end{aligned}$$

while in the wave region we have

$$\varPsi=\varPsi^{(0)}+\mathscr{M}\varPsi^{(1)}+\mathscr{M}^2\varPsi^{(2)}+\dots. \qquad (16)$$

We restrict our analysis to order $\mathcal{O}(\mathscr{M}^2)$, because at higher orders the calculations become very involved.

A. The solution to $\mathcal{O}(\mathscr{M}^0)$

At this order, the solution in the wave region is determined by the ordinary wave equation $(\boldsymbol{\nabla}^2+k^2)\varPsi^{(0)}=0$, where the nondimensional wave number k is defined as $k=\varOmega$. We expand the general solution to zeroth order, corresponding to our incoming wave, in the following way into partial waves of order ℓ:

$$\varPsi^{(0)}=\sum_{\ell=-\infty}^{\infty}A_\ell\,J_{|\ell|}(kR)\,e^{i(\ell\phi-\varOmega t)}, \qquad (17)$$

where the $J_{|\ell|}$ are Bessel functions. The form of $\varPsi^{(0)}$ in the vortical region may be determined by expanding the Bessel functions for small R, yielding

$$\begin{aligned}\varPsi^{(0)} \;=\;& A_0e^{-i\varOmega t}+\mathscr{M}kr\sum_{\ell=\pm1}\frac{A_\ell}{2}e^{i(\ell\phi-\varOmega t)} \\ &+\mathscr{M}^2\frac{k^2r^2}{4}\left(\sum_{\ell=\pm2}\frac{A_\ell}{2}e^{i(\ell\phi-\varOmega t)}-A_0e^{-i\varOmega t}\right) \\ &+\mathcal{O}(\mathscr{M}^3). \qquad (18)\end{aligned}$$

This will be needed later to carry out the matching.

In the vortical region, Eq. (12) to this order gives the ordinary Poisson equation $\boldsymbol{\nabla}^2(\psi^{(0)}+\eta^{(0)})=0$, with the following general solution

$$\psi^{(0)}+\eta^{(0)}=\sum_{\ell=-\infty}^{\infty}a_\ell\,(kr)^{|\ell|}\,e^{i(\ell\phi-\varOmega t)}. \qquad (19)$$

In the wave region, $r^{|\ell|}=\mathcal{O}(\mathscr{M}^{-|\ell|})$, ruling out the existence of terms with $|\ell|>0$, since these would be of *negative* order in the Mach number. Thus, $\psi^{(0)}+\eta^{(0)}=a_0$.

A decomposition of $\zeta^{(0)}$ into azimuthal modes according to $\zeta^{(0)}=\sum_{\ell=-\infty}^{\infty}\zeta_\ell^{(0)}(r)\,e^{i(\ell\phi-\varOmega t)}$ reduces Eq. (14) to

$$\left[\frac{\mathrm{d}^2}{\mathrm{d}r^2}+\frac{1}{r}\frac{\mathrm{d}}{\mathrm{d}r}-\frac{\ell^2}{r^2}+\frac{\ell\omega_0'}{r\varOmega-\ell v_0}\right]\zeta_\ell^{(0)}(r)=0. \qquad (20)$$

We restrict ourselves to the case where there are no critical layers present, $\varOmega\neq\ell v_0(r)/r$ for all ℓ, r [19]. In this case, the solutions of this radial Rayleigh equation are known to behave for large r as $\zeta_\ell^{(0)}(r)\to\alpha_\ell\left(r^{|\ell|}+\beta_\ell r^{-|\ell|}\right)$, immediately giving $\alpha_\ell=0$ for all ℓ. Furthermore, using identity (13) and $v_1^{(0)}=\boldsymbol{\nabla}\psi^{(0)}+\boldsymbol{\xi}^{(0)}=0$ as well as $\boldsymbol{\nabla}\psi^{(0)}=-\boldsymbol{\nabla}\eta^{(0)}$, we can derive the equation $-i\Omega\eta^{(0)}+\frac{v_0}{r}\frac{\partial\eta^{(0)}}{\partial\phi}=$ const., from which it follows that $\eta^{(0)}=Ce^{i(\Omega r/v_0)\phi}+$ const. Therefore, as $\eta^{(0)}$ should be uniquely defined at each point in space, $C{=}0$ in the presently considered case; $\eta^{(0)}$ thus equals a constant, which is zero because it must vanish at large r.

Matching the solution in the vortical region with the wave region gives $a_0=A_0$, thus

$$\psi^{(0)}=A_0\,e^{-i\varOmega t}, \quad \eta^{(0)}=\zeta^{(0)}=0. \qquad (21)$$

To this lowest (zeroth) order in the Mach number, there is thus no perturbative flow in the vortical region, but since $\psi^{(0)}\neq0$, there are density perturbations present.

B. The solution to $\mathcal{O}(\mathscr{M})$

In the wave region, we have once more the ordinary wave equation $(\boldsymbol{\nabla}^2+k^2)\varPsi^{(1)}=0$. We require the solution to this equation to be causal; therefore we write

$$\varPsi^{(1)}=\sum_{\ell=-\infty}^{\infty}B_\ell H_\ell^{(1)}(kR)\,e^{i(\ell\phi-\varOmega t)}, \qquad (22)$$

where the Hankel functions of the first kind $H_\ell^{(1)}$ correspond to outgoing cylindrical waves. Due to the irregular behavior of the Hankel functions at the origin, none of the terms in the above sum matches to the solution in the vortical region, as becomes clear from Eq. (26) below; therefore, $B_\ell = 0$. Thus, $\Psi^{(1)} = 0$ in the wave region, reflecting the simple fact that there is no sound radiation to linear order in the Mach number into the wave region.

In the vortical region, $\psi^{(1)} + \eta^{(1)}$ is once more determined by the ordinary Poisson equation, giving $\psi^{(1)} + \eta^{(1)} = \sum_{\ell=-1}^{1} b_\ell \,(kr)^{|\ell|}\, e^{i(\ell\phi - \Omega t)}$, where the sum is limited by the fact that higher-order terms would be of negative order in the Mach number. The quantity $\zeta^{(1)}$ ist determined by the following equation:

$$\left[\left(-i\Omega + \frac{v_0}{r}\frac{\partial}{\partial\phi}\right)\boldsymbol{\nabla}^2 - \frac{\omega_0'}{r}\frac{\partial}{\partial\phi}\right]\zeta^{(1)} = -k\omega_0' \sum_{\ell=\pm1} b_\ell e^{i(\ell\phi - \Omega t)}. \tag{23}$$

Using the ansatz $\zeta_p^{(1)} = f(r)\sum_{\ell=\pm1} b_\ell e^{i(\ell\phi-\Omega t)}$ and $\boldsymbol{\nabla}^2 v_0 = \partial_r(r^{-1}\partial_r(r v_0)) = \omega_0'$, we immediately see that the equation above is solved once one chooses $f = -i v_0$. The asymptotics of the solution thus obtained for $r \to \infty$ is given by

$$\zeta_{\mathrm{p}}^{(1)} \to -i\frac{\gamma}{r}\sum_{\ell=\pm1} b_\ell e^{i(\ell\phi-\Omega t)}. \tag{24}$$

One gets the full solution to $\zeta^{(1)}$ by adding the general solution $\zeta_{\mathrm{h}}^{(1)}$ of the homogeneous Rayleigh equation to $\zeta_{\mathrm{p}}^{(1)}$. Since $\zeta_{\mathrm{p}}^{(1)} \to 0$ for $r \to \infty$, $\zeta_{\mathrm{h}}^{(1)} \to 0$ for $r \to \infty$ as well. Using the homogeneous version of Eq. (23), which can be transformed to a Rayleigh equation like Eq. (20); the same argument as given after Eq. (20) then yields $\zeta_{\mathrm{h}}^{(1)} = 0$. Since we have

$$\begin{aligned}\boldsymbol{e}_z \times \boldsymbol{\nabla}\zeta^{(1)} &\to i\frac{\gamma}{r^2}\sum_{\ell=\pm1} b_\ell(\boldsymbol{e}_\phi + i\ell\boldsymbol{e}_r)e^{i(\ell\phi-\Omega t)} \\ &= \boldsymbol{\nabla}\left[\frac{\gamma}{r}\sum_{\ell=\pm1} \ell\, b_\ell\, e^{i(\ell\phi-\Omega t)}\right]\end{aligned} \tag{25}$$

for $r \to \infty$, the asymptotics of $\eta^{(1)}$ for $r \to \infty$ can be determined to be $\eta^{(1)} \to -\frac{\gamma}{r}\sum_{\ell=\pm1} \ell\, b_\ell\, e^{i(\ell\phi-\Omega t)}$, thus giving

$$\psi^{(1)} \to b_0 + \sum_{\ell=\pm1} b_\ell\left[kr + \ell\frac{\gamma}{r}\right] e^{i(\ell\phi-\Omega t)} \tag{26}$$

for $r \to \infty$. By comparison with the solution in the wave region, we obtain $b_0 = 0$ and $b_{\pm1} = A_{\pm1}/2$.

The rotational part of $\boldsymbol{v}_1^{(1)}$, $\boldsymbol{e}_z \times \boldsymbol{\nabla}\zeta^{(1)}$, describes the oscillatory motion of the vortex core induced by the sound wave [20], which may be seen as follows. Near the origin, we may expand the incoming wave into a plane-wave form $\Psi^{(0)} \approx \psi_0 e^{i(\boldsymbol{k}\cdot\boldsymbol{R}-\Omega t)}$, for which the expansion coefficients are given by $A_\ell = \psi_0 e^{-i\ell\phi_k}\, i^{|\ell|}$, with ϕ_k the angle which the propagation vector $\boldsymbol{k} = k(\cos\phi_k, \sin\phi_k, 0)$ makes with the x axis. We therefore have $\boldsymbol{e}_z \times \boldsymbol{\nabla}\zeta^{(1)} \approx \boldsymbol{e}_z \times (\frac{1}{2}\psi_0\omega_0\boldsymbol{e}_k)e^{-i\Omega t}$. Then, using $\boldsymbol{v}_0 \approx \frac{1}{2}(\boldsymbol{\omega}_0 \times \boldsymbol{r})$ near the origin, $\boldsymbol{e}_z \times (\frac{1}{2}\psi_0\omega_0\boldsymbol{e}_k)e^{-i\Omega t} \approx -\psi_0(e^{-i\Omega t}\boldsymbol{e}_k \cdot \boldsymbol{\nabla})\boldsymbol{v}_0$. We see that the term $\boldsymbol{e}_z \times \boldsymbol{\nabla}\zeta^{(1)}$ causes a displacement $\boldsymbol{r}_0(t) \equiv \psi_0 e^{-i\Omega t}\boldsymbol{e}_k$ of the vortex line. It follows for the resulting displacement velocity $\frac{\partial}{\partial t}\boldsymbol{r}_0(t) = -i\Omega\psi_0 e^{-i\Omega t}\boldsymbol{e}_k = \boldsymbol{\nabla}\Psi^{(0)}|_{\boldsymbol{r}=0}$, so that the vortex indeed moves with the velocity in the incident sound wave to this order in the Mach number.

C. The solution to $\mathcal{O}(\mathscr{M}^2)$

In the vortical region, $\boldsymbol{\nabla}^2\left(\psi^{(2)} + \eta^{(2)}\right) = -k^2 A_0 e^{-i\Omega t}$, with the following solution:

$$\psi^{(2)} + \eta^{(2)} = -\frac{A_0 e^{-i\Omega t}}{4}k^2 r^2 + \sum_{\ell=-2}^{2} c_\ell\,(kr)^{|\ell|}\, e^{i(\ell\phi-\Omega t)}. \tag{27}$$

Comparison with (18) leads to $c_0 = c_{\pm1} = 0$ and $c_{\pm2} = A_{\pm2}/8$. In the wave region, $\Psi^{(2)}$ obeys a forced wave equation:

$$(\boldsymbol{\nabla}^2 + k^2)\Psi^{(2)} = -2ik\gamma R^{-2}\partial_\phi\Psi^{(0)}. \tag{28}$$

In the appendix, it is shown that the solution to this equation may be written as a contour integral. The result is (with $\mathcal{C}$ denoting the integration contour shown in the appendix):

$$\Psi_{\mathrm{p}}^{(2)} = \mathrm{sgn}(\ell)\, i\frac{k\gamma}{2\pi}A_\ell \sum_{\ell=-\infty}^{\infty} \Phi_\ell(R)e^{i(\ell\phi-\Omega t)} \tag{29}$$

where the expansion coefficients are given by

$$\Phi_\ell(R) = \int_{\mathcal{C}}\left(\ln t + i\frac{\pi}{2}\right)\exp\left[i\frac{kR}{2}\left(t - \frac{1}{t}\right)\right]\frac{\mathrm{d}t}{t^{|\ell|+1}}. \tag{30}$$

We can write the general solution to Eq. (28) as $\Psi^{(2)} = \Psi_{\mathrm{p}}^{(2)} + \Psi_{\mathrm{h}}^{(2)}$, with $\Psi_{\mathrm{h}}^{(2)}$ being the general solution of the homogeneous equation. For large R, $\Psi_{\mathrm{p}}^{(2)}$ corresponds to an outgoing wave (cf. the appendix). Therefore, $\Psi_{\mathrm{h}}^{(2)}$ should also contain only the outgoing wave:

$$\Psi_{\mathrm{h}}^{(2)} = \sum_{\ell=-\infty}^{\infty} C_\ell\, H_\ell^{(1)}(kR)e^{i(\ell\phi-\Omega t)}. \tag{31}$$

The remaining coefficients are determined by matching to the solution in the vortical region, giving $C_\ell = 0$ for $\ell \neq \pm1$ and

$$C_{\pm1} = \pm i\frac{\pi}{4}k\gamma A_{\pm1}. \tag{32}$$

The solution to quadratic order in the Mach number, $\varPsi^{(2)}$, is thus made up of two parts. The first part, $\varPsi_{\rm p}^{(2)}$, may be interpreted to represent the interaction of the incident wave with the long-range velocity field of the vortex, while the second, $\varPsi_{\rm h}^{(2)}$ stands for the interaction of the wave with the vortex core [19].

IV. FORCE ON THE VORTEX FROM PHONON SCATTERING

In this section we will return to the original (dimensional) variables of section II. Far from the vortex, our solution takes on the following asymptotical form: $\psi = \sum_\ell \psi_\ell(r)\, e^{i(\ell\phi - \Omega t)}$ where, using the asymptotic form of the expansion coefficients $\varPhi_\ell(R)$ in the far-field solution (29), derived in the appendix [Eq. (A5)], we have

$$\psi_\ell = \frac{e^{-i\frac{\pi}{2}|\ell| - i\frac{\pi}{4}}}{\sqrt{2\pi k r}} A_\ell \left\{ (1 + 2i\delta_\ell) e^{ikr} + e^{-ikr + i\pi|\ell| + i\frac{\pi}{2}} \right\}. \tag{33}$$

Here, the phase shifts δ_ℓ are given by

$$\delta_\ell = \begin{cases} \mp\frac{\pi}{4}\frac{k\gamma}{c_\infty} & ; \ell = \pm 1, \\ -\mathrm{sgn}(\ell)\frac{\pi}{2}\frac{k\gamma}{c_\infty} & ; \ell \neq \pm 1, \end{cases} \tag{34}$$

which are idential to those obtained by Sonin [4]. To evaluate the momentum transfer between sound wave and vortex, we consider the time-averaged momentum-flux tensor, which reads [4, 8, 14]:

$$\langle \varPi_{ij} \rangle = \left(\frac{c_\infty^2}{\rho_\infty} \frac{\langle \rho_1^2 \rangle}{2} - \rho_\infty \frac{\langle v_1^2 \rangle}{2} \right) \delta_{ij} + \rho_\infty \langle v_{1i} v_{1j} \rangle, \tag{35}$$

where ρ_∞ is the constant density far from the vortex. We have to insert into this expression the time-averaged expressions for the density and the velocity expressed in terms of ψ and $\boldsymbol{\xi}$ according to (4). Since we are looking at the region far from the vortex, we may use the simplified expressions:

$$\rho_1 = -\frac{\rho_\infty}{c_\infty^2} \mathrm{Re}\,[\partial_t \psi] = \frac{i\rho_\infty k}{2c_\infty}(\psi^* - \psi), \tag{36}$$

$$v_{1i} = \mathrm{Re}\,[\partial_i \psi] = \frac{1}{2}(\partial_i \psi^* + \partial_i \psi), \tag{37}$$

which, when inserted into the time-averaged momentum-flux tensor (35), yield

$$\langle \varPi_{ij} \rangle = \frac{\rho_\infty}{4} [\partial_i \psi^* \partial_j \psi + \partial_i \psi \partial_j \psi^* + (k^2 |\psi|^2 - |\boldsymbol{\nabla} \psi|^2) \delta_{ij}]. \tag{38}$$

This comprises a general expression for the time-averaged momentum-flux tensor in the far-region where $\boldsymbol{v}_0 \approx 0$.

The projection F_{ϕ_0} of the force on the vortex per unit length $\boldsymbol{F}$ in the direction of the unit vector $\boldsymbol{n}_0 = (\cos\phi_0, \sin\phi_0, 0)$ is given by

$$F_{\phi_0} = \langle \boldsymbol{F} \cdot \boldsymbol{n}_0 \rangle = -\oint \langle \varPi_{ij} \rangle n_i \, \mathrm{d}a_j. \tag{39}$$

Inserting (38) into the above expression, and choosing a cylindrical surface around the vortex as our contour of integration, we get

$$F_{\phi_0} = -\frac{\rho_\infty}{4} \int_{-\pi}^{\pi} \mathrm{d}\phi\, r \cos(\phi - \phi_0)\, (\partial_r \psi^* \partial_r \psi + k^2 \psi^* \psi). \tag{40}$$

Making use of (33), and performing the angular integration, we find

$$F_{\phi_0} = \rho_\infty k \sum_{\ell=-\infty}^{\infty} \mathrm{Re} \left[i^{|\ell|+1-|\ell+1|} e^{i\phi_0} A_\ell^* A_{\ell+1} (\delta_\ell - \delta_{\ell+1}) \right]. \tag{41}$$

Employing the obtained phase shifts for sound-vortex scattering, Eq. (34), we arrive at the following final expression for the force:

$$F_{\phi_0} = \frac{\rho_\infty k^2}{c_\infty} \frac{\pi}{4} \gamma\, \mathrm{Re} \left[e^{i\phi_0} \sum_{\ell=0}^{1} (A_\ell^* A_{\ell+1} - A_{-\ell-1}^* A_{-\ell}) \right]. \tag{42}$$

The above equation represents the general result for the force on the vortex from phonon scattering, containing partial waves up to order $|\ell| = 2$, valid for an arbitrary incident wave form.

Note that there are no singularities in (42) present due to partial wave summation, because we have assumed an arbitrary incoming wave form right from the start. Only now, *after* obtaining the general result for the force in terms of the expansion coefficients A_ℓ, we use a plane wave approximation for the incoming wave near the origin. Then, with $A_\ell = \psi_0 e^{-i\ell\phi_k}\, i^{|\ell|}$,

$$F_{\phi_0} = -\frac{\rho_\infty k^2}{c_\infty} \pi\gamma |\psi_0|^2 \sin(\phi_0 - \phi_k). \tag{43}$$

Using the following expression for the momentum density of a plane sound wave, $\boldsymbol{j} = |\psi_0|^2 \frac{\rho_\infty k}{2c_\infty} \boldsymbol{k}$ [8], Eq. (43) may easily be generalized to yield the parallel and transverse components of the force,

$$\boldsymbol{F}_{||} = 0, \quad \boldsymbol{F}_\perp = -\boldsymbol{\varGamma} \times \boldsymbol{j}, \tag{44}$$

where $\boldsymbol{\varGamma} = 2\pi\gamma \boldsymbol{e}_z$ and $\boldsymbol{j}$ is the momentum density of the wave at the location of the vortex. The coordinate invariant form of (44) is obviously not restricted to the case where the vortex axis points into the z-direction. For a vortex in a superfluid we have to replace $\boldsymbol{j}$ by the thermally averaged momentum density of the phonons $\boldsymbol{j}_{\rm ph}$, which in terms of the quantities of two-fluid hydrodynamics reads $\boldsymbol{j}_{\rm ph} = \rho_n(\boldsymbol{v}_n - \boldsymbol{v}_s)$, where $\boldsymbol{v}_n$ and $\boldsymbol{v}_s$ are normal and superflow velocities, respectively. For a vortex which is at rest with respect to the superfluid condensate ($\boldsymbol{v}_L = \boldsymbol{v}_s$) this gives

$$\boldsymbol{F}_{\rm Iordanskii} = \rho_n [\boldsymbol{\varGamma} \times (\boldsymbol{v}_L - \boldsymbol{v}_n)], \tag{45}$$

which is the Iordanskii force. We remark that the arguments presented in [8] suggest that the above form of the force remains invariant (to the considered order in the Mach number), also if the vortex moves with respect to the superflow.

V. CONCLUSION

We have solved the problem of scattering of sound waves by a vortex using an expansion into partial waves and the method of matched asymptotic expansions, up to the relevant quadratic order in the Mach number. To the best of our knowledge, the analytical form of the solution for the scattered wave in the form of a Schläfli contour integral, Eq. (29), has not been given yet in the literature. By evaluating the momentum-flux tensor far from the vortex, we derived a general expression for the force exerted by the sound wave on the vortex. The force depends only on the lowest partial waves ($|\ell| \leq 2$) and is thus completely determined by the shape of the incident wave in a small region around the vortex axis with a diameter of the order of a few wavelengths (an analogous result was obtained by Shelankov for an incident wave of beam-shaped form [6]). We demonstrated that the usage of a "physical" incident wave of arbitrary shape (and finite extent) removes the mathematical difficulties which plagued previous work employing the partial wave method, where the total transverse cross section, obtained after performing a summation over all partial waves, contained singularities (was not convergent) [2, 4, 5]. The resulting global, singularity-free solution of the scattering problem we presented then leads, in conjunction with conventional two-fluid hydrodynamics, to the Iordanskii force on the vortex, confirming its existence as a result of the asymmetrical scattering of sound waves at a vortex.

Direct experimental confirmation of the Iordanskii force has so far been elusive. A conceivable experimental procedure to verify the existence of the Iordanskii force in a superfluid is to study the influence of sound pulses in Bose-Einstein condensates, created by Bragg scattering [21], on the vortex. Using Bose-Einstein condensates has, inter alia, the advantage of being able to create sound wave pulses in an accurately controlled manner, with variable large momenta of the order of c/γ. We point out that our approach predicts the transverse force correctly also if the incoming wave is not a plane wave, which will in general be the case in the inhomogeneously trapped condensates. If the vortex is pinned by an optically created external potential to fix it in one position, and, shortly after the sound wave has been created, the pinning potential is turned off, one should in principle be able to observe the transverse displacement of the vortex due to the sound wave scattered by it.

APPENDIX A: FORCED WAVE EQUATION

In this appendix, we derive the solution (29) to the forced wave equation (28). Employing the ansatz (29), Eq. (28) reduces to

$$\left[R^2\frac{\mathrm{d}^2}{\mathrm{d}R^2}+R\frac{\mathrm{d}}{\mathrm{d}R}+k^2R^2-\ell^2\right]\mathit{\Phi}_\ell=-4|\ell|\pi i\,J_{|\ell|}(kR). \tag{A1}$$

We write the solution to the above inhomogeneous Bessel equation as a contour integral. On the right-hand side, using the Schläfli integral representation for the Bessel functions [22], with the contour $\mathcal{C}$ shown in Fig. 1, we express $J_{|\ell|}(kR)$ as

$$J_{|\ell|}(kR)=\frac{1}{2\pi i}\int_{\mathcal{C}}\exp\left[\frac{kR}{2}\left(t-\frac{1}{t}\right)\right]\frac{\mathrm{d}t}{t^{|\ell|+1}}. \tag{A2}$$

On the left-hand side, inspired by the above Schläfli integral representation, we try the following ansatz:

$$\mathit{\Phi}_\ell=\int_{\mathcal{C}}g_\ell(t)\,\exp\left[\frac{kR}{2}\left(t-\frac{1}{t}\right)\right]\frac{\mathrm{d}t}{t^{|\ell|+1}}. \tag{A3}$$

After partially integrating the left-hand side and using the well known recursion relation $J_{\ell-1}(x)+J_{\ell+1}(x)=\frac{2\ell}{x}J_\ell(x)$ for the Bessel functions on the right-hand side, we find that the function $g_\ell(t)$ is determined by $\frac{dg_\ell}{dt}=t^{-1}$. If we choose the integration constant such that

$$g_\ell(t)=\ln t+i\frac{\pi}{2}, \tag{A4}$$

then the $\mathit{\Phi}_\ell$ correspond to causal solutions of (A1). This may easily be verified by applying the method of steepest descent, yielding the following asymptotics:

$$\mathit{\Phi}_\ell\to-2\pi\sqrt{\frac{\pi}{2kR}}\exp\left[i\left(kR-|\ell|\frac{\pi}{2}-\frac{\pi}{4}\right)\right] \tag{A5}$$

for $R\to\infty$. Notice that $\mathit{\Psi}_\mathrm{p}^{(2)}$ is regular at the origin, $\mathit{\Psi}_\mathrm{p}^{(2)}\to 0$ as $R\to 0$, as can easily be confirmed by expanding the integrand for small R and integrating.



FIG. 1: Schläfli contour of integration

---

[1] S. V. Iordanskii, Sov. Phys. JETP **22**, 160 (1966).

[2] D. J. Thouless, P. Ao, and Q. Niu, Phys. Rev. Lett. **76**, 3758 (1996).

[3] G. E. Volovik, JETP Lett. **62**, 65 (1995); Phys. Rev. Lett. **77**, 4687 (1996).

[4] E. B. Sonin, Phys. Rev. B **55**, 485 (1997); Phys. Rev. Lett. **81**, 4276 (1998).

[5] C. Wexler, Phys. Rev. Lett. **79**, 1321 (1997); C. Wexler and D. J. Thouless, Phys. Rev. B **58**, R8897 (1998).

[6] A. L. Shelankov, Europhys. Lett. **43**, 623 (1998).

[7] G. E. Volovik, JETP Lett. **67**, 881 (1998).

[8] M. Stone, Phys. Rev. B **61**, 11780 (2000).

[9] D. J. Thouless, M. R. Geller, W. F. Vinen, J.-Y. Fortin, and S. W. Rhee, Phys. Rev. B **63**, 224504 (2001).

[10] E. B. Sonin, preprint cond-mat/0104221, published in *Vortices in Unconventional Superconductors and Superfluids*, R. P. Hübener, N. Schopohl, and G. E. Volovik (Eds.), Springer, Berlin 2002, pp. 119.

[11] C. Zhang, A. M. Dudarev, and Q. Niu, Phys. Rev. Lett. **97**, 040401 (2006).

[12] L. P. Pitaevskii, Sov. Phys. JETP **8**, 888 (1959).

[13] A. L. Fetter, Phys. Rev. **136**, A1488 (1964).

[14] M. Stone, Phys. Rev. E **62**, 1341 (2000).

[15] S. E. P. Bergliaffa, K. Hibberd, M. Stone, and M. Visser, Physica D **191**, 121 (2004).

[16] W. G. Unruh, Phys. Rev. Lett. **46**, 1351 (1981).

[17] A. D. Pierce, J. Acoust. Soc. Am. **87**, 2292 (1990).

[18] M. V. Fedoryuk (Ed.), *Partial Differential Equations V*, Springer, Berlin 1998.

[19] R. Ford und S. G. Llewellyn Smith, J. Fluid. Mech. **386**, 305 (1999).

[20] S. Leblanc, J. Fluid. Mech. **414**, 315 (2000).

[21] D. M. Stamper-Kurn, A. P. Chikkatur, A. Görlitz, S. Inouye, S. Gupta, D. E. Pritchard, and W. Ketterle, Phys. Rev. Lett. **83**, 2876 (1999).

[22] L. Schläfli, Math. Ann. **3**, 134 (1871).